

Introduction to Pediatric Dermatology

Bernard A. Cohen

ANATOMY AND FUNCTION OF THE SKIN

Most of us think of skin as a simple, durable covering for the skeleton and internal organs. Yet skin is actually a very complex and dynamic organ consisting of many parts and appendages (Fig. 1.1). The outermost layer of the epidermis, the stratum corneum, is an effective barrier to the penetration of irritants, toxins, and organisms, as well as a membrane that holds in body fluids. The remainder of the epidermis, the stratum granulosum, stratum spinosum, and stratum basale, manufactures this protective layer. Melanocytes within the epidermis are important for protection against the harmful effects of ultraviolet light, and the Langerhans cells and other dendritic cells are one of the body's first lines of immunologic defense and play a key role in systemic and cutaneous diseases such as drug reactions and infections.

The dermis, consisting largely of fibroblasts and collagen, is a tough, leathery, mechanical barrier against cuts, bites, and bruises. Its collagenous matrix also provides structural support for a number of cutaneous appendages. Hair, which grows from follicles deep within the dermis, is important for cosmesis, as well as protection from sunlight and particulate matter. Sebaceous glands arise as an outgrowth of the hair follicles. Oil produced by these glands helps to lubricate the skin and contributes to the protective function of the epidermal barrier. The nails are specialized organs of manipulation that also protect sensitive digits. Thermoregulation of the skin is accomplished by eccrine sweat glands and changes in the cutaneous blood flow regulated by glomus cells. The skin also contains specialized receptors for heat, pain, touch, and pressure. Sensory input from these structures helps to protect the skin surface against environmental trauma. Beneath the dermis, in the subcutaneous tissue, fat is stored as a source of energy and also acts as a soft protective cushion.

EXAMINATION AND ASSESSMENT OF THE SKIN

The skin is the largest and most accessible and easily examined organ of the body, and it is often the organ of most frequent concern and quality of life for the patient. Therefore all practitioners should be able to recognize basic skin diseases and dermatologic clues to systemic disease.

Optimal examination of the skin is best achieved in a well-lit room. The clinician should inspect the entire skin surface,

including the hair, nails, scalp, and mucous membranes. This may present particular problems in infants and teenagers, because it may be necessary to examine the skin in small segments to prevent cooling or embarrassment, respectively. Although no special equipment is required, a hand lens and side lighting are useful aids in the assessment of skin texture and small, discrete lesions. In many offices, the otoscope can be adapted for this purpose by removing the plastic speculum.

There are also a number of relatively inexpensive portable dermatoscopic devices, which can also be used to enhance the examination (also known as epiluminescence microscopy). These instruments have traditionally provided a magnified ($\times 10$) view of the skin with a nonpolarized light source, a transparent plate, and a liquid medium between the dermatoscope and the skin. This allows for a view of the superficial structures in the skin without interference from surface reflections. Dermatoscopic heads can be purchased for otoscope/ophthalmoscope handpieces, and mineral oil or alcohol gel can be applied directly to the skin lesion. Newer devices allow for toggling between nonpolarized and polarized light, which provides visualization of deeper structures in the dermis. (Fig. 1.2).

Despite the myriad of conditions affecting skin, a systematic approach to the evaluation of a rash facilitates and simplifies the process of developing a manageable differential diagnosis. After assessing the general health of a child, the practitioner should obtain a detailed history of the cutaneous symptoms, including the date of onset, inciting factors, evolution of lesions, and presence or absence of pruritus. Recent immunizations, infections, drugs, and allergies may be directly related to new rashes. The family history may suggest a hereditary or contagious process, and the clinician may need to examine other members of the family. A review of nursery records and images provided by parents and in the electronic record will help document the presence and evolution of congenital and acquired lesions.

Attention should then turn to the distribution and pattern of the rash. The distribution refers to the location of the skin findings, while the pattern defines a specific anatomic or physiologic arrangement. For example, the distribution of a rash may include the extremities, face, or trunk, while the pattern could be flexural or intertriginous areas (Fig. 1.3a). Other common patterns include sun-exposed sites, acrodermatitis (predilection for the distal extremities), pityriasis rosea (truncal, following the skin cleavage lines), clothing-protected sites, acneiform rashes, lines of Blaschko, and segmental lesions (Fig. 1.3b–h).




Fig. 1.1 (a) Skin photomicrograph and (b) schematic diagram of normal skin anatomy.




Fig. 1.2 Examination of pigmented nevi with (a) handheld DermLite dermatoscope by 3Gen, (b) DermLite DL4W dermatoscope by 3Gen, and (c) Welch Allyn otoscope.

Next, the clinician should consider the local organization and configuration of the lesions, defining the relationship of primary and secondary lesions to one another in a given location (Table 1.1) and the shape of the lesions. Are the lesions diffusely scattered or clustered (herpetiform)? Are they dermatomal, linear, serpiginous, circular, annular, or reticulated?

The depth of the lesions in the skin, as noted by both observation and palpation, may also give further clues (Table 1.2). Disruption of the normal skin markings by scale, papules, vesicles, or pustules points to the involvement of the epidermis. Alterations in skin color alone can occur in epidermal and dermal processes. In disorders of pigmentation, the color of the pigment may suggest the anatomic depth of the lesion. Shades of brown are present in flat junctional nevi, lentiginous,

and café-au-lait spots, where the increased pigment resides in the epidermis or superficial dermis. In Mongolian spots and nevus of Ota, the Tyndall effect results in bluish-green to gray macules from melanin uniformly distributed in the mid-dermis. If the epidermal markings are normal but the lesion is elevated, the disorder usually involves the dermis. Dermal lesions have well-demarcated firm borders. Nodules and tumors deep in the dermis or subcutaneous tissue can distort the surface markings, which are otherwise intact. Some deep-seated lesions can only be appreciated by careful palpation.

Lesion color can provide important clues for diagnosis and the pathophysiology of the underlying process (Table 1.3). Brown, blue, gray, bronze, and black lesions are associated with disorders that alter normal pigmentation, while white lesions








Fig. 1.3 (a–h) Pattern diagnosis.

TABLE 1.1 Organization and Configuration of Lesions

Linear	Dermatomal	Serpiginous	Annular	Herpetiform	Reticulated	Filiform (Thread-like)	Geographic
Epidermal nevi	Herpes zoster	Psoriasis	Ringworm	Herpes simplex infection	Cutis marmorata	Wart	Psoriasis
Lichen striatus	Vitiligo	Erythema marginatum	Granuloma annulare	Herpes zoster	Livedo reticularis	Dermatosis papulosa nigra	Geographic tongue
Contact dermatitis	Nevus depigmentosus	Cutaneous larvae migrans	Subacute cutaneous lupus	Dermatitis herpetiformis	Congenital phlebectasia	Syringocystadenoma papilliferum	Nummular eczema
Warts	Becker nevus	Elastosis perforans serpiginosa	Atopic dermatitis		Reticulated and confluent papillomatosis	Skin tag	Erythema annulare centrifugum
Ichthyosis	Café-au-lait spot		Erythema annulare centrifugum		Erythema ab igne	Pigmented nevus	
Psoriasis	Port-wine stain		Erythema chronicum migrans				
Porokeratosis			Erythema marginatum				
Incontinentia pigmenti							

TABLE 1.2 Anatomic Depth of Lesions

Cutaneous Structure	Physical Findings	Specific Skin Disorder	
Epidermis	Altered surface markings Scale, vesicle, crust Color changes (black, brown, white)	Impetigo Café-au-lait spot Atopic dermatitis Vitiligo Freckle	
Epidermis + dermis	Altered surface markings Scale, vesicle, crust Distinct borders Color changes (black, brown, white, and/or red) Edema	Psoriasis Atopic dermatitis Cutaneous lupus erythematosus	
Dermis	Normal surface markings Color changes Altered dermal firmness	Urticaria Granuloma annulare Hemangioma Blue nevus	
Subcutaneous tissue	Normal surface markings Normal or red skin color Altered skin firmness	Hematoma Cold panniculitis Erythema nodosum	

may be associated with loss of normal pigmentation or the accumulation of scale, crust, or exudates. Red and blue lesions are associated with inflammatory and vascular processes. Non-blanching blue or purple lesions should suggest the presence of purpura. Yellow lesions occur when the skin is infiltrated with inflammatory or tumor cells containing lipid. Other pigments from topical agents (e.g. silver, gold), oral medications (e.g. minocycline, amiodarone), foreign bodies (e.g. asphalt, tattoo

pigments), and infectious agents (e.g. *Pseudomonas* species, *Corynebacterium* species) may impart specific colors to cutaneous lesions.

Finally, the clinician may develop a differential diagnosis using the morphology of the cutaneous lesions. Primary lesions (macules, papules, plaques, vesicles, bullae, pustules, wheals, nodules, and tumors) arise *de novo* in the skin (Fig. 1.4). Secondary lesions (scale, crust, erosions, ulcers, scars with atrophy

TABLE 1.3 Lesion Color

Red	Purple	Brown	Gray	Blue	Bronze	Green	Yellow
 Inflammatory disorders, such as eczema, psoriasis, urticaria, erythema chronicum migrans, and other figurate erythemas	 Purpura, vascular malformations, hemangiomas, hematoma	 Pigmented nevus, post-inflammatory hyperpigmentation, lentigo, ephelid (freckle), café-au-lait spot, epidermal nevus	 Mongolian spot, graphite tattoo, nevus of Ota	 Tattoo, vascular malformation, hemangiomas, blue nevus, Mongolian spot	 Progressive-pigmented purpuric dermatosis, resolving hematoma, phyto-photodermatitis	 Tattoo, pseudomonas infection, deposition of minocycline, Mongolian spot, resolving hematoma	 Xantho-granuloma, xanthoma, sebaceous hyperplasia, epidermal inclusion cyst

and/or fibrosis, excoriations, and fissures) evolve from primary lesions or result from scratching of primary lesions by the patient (Fig. 1.5).

The practitioner who becomes comfortable with dermatology will integrate all of these approaches into their evaluation of a child with a skin problem. This will be reflected in the clinically focused format of this text.

Each chapter will finish with an algorithm that summarizes the material in a differential diagnostic flow pattern. The limited bibliography includes comprehensive, historically significant, and/or well-organized reviews of the subject. Readers may also find some of the texts and online further reading listed at the end of this chapter useful.

DIAGNOSTIC TECHNIQUES

Potassium Hydroxide Preparation

There are a number of rapid, bedside diagnostic procedures in dermatology. One of the most useful techniques is a wet mount of skin scrapings for microscopic examination (Fig. 1.6). Potassium hydroxide (KOH) 20% is used to change the optic properties of skin samples and make scales more transparent. The technique requires practice and patience.

The first step is to obtain the material by scraping loose scales at the margin of a lesion, nail parings, subungual debris, or the small, pearly globules from a molluscum body. Short residual hair stubs (black dots in tinea capitis) may also be painlessly shaved off the scalp with a #15 blade. Scale is placed on the slide and moved to the center with a cover slip. One or two drops of KOH are added and gently warmed with a match or the microscope light. Boiling the specimen will introduce artifacts and should be avoided, so sitting the specimen aside for 5 min is an alternative to gentle heating. Excess KOH can be removed with a paper towel applied to the edge of the cover slip. Thick specimens may be more easily viewed after gentle but firm pressure is applied to the cover slip with a pencil eraser. Thick scale will also dissolve after being set aside for 15–20 min.

View the preparation under a microscope, with the condenser and light at low levels to maximize contrast, and with the

objective at $\times 10$. Focus up and down as the entire slide is rapidly scanned. True hyphae are long branching green hyaline rods of uniform width that cross the borders of epidermal cells. They often contain septae. False positives may be vegetative fibers, cell borders, or other artifacts. Yeast infections show budding yeast and pseudohyphae. Molluscum bodies are oval discs that have homogeneous cytoplasm and are slightly larger than keratinocytes. In hair fragments, the fungi appear as small, round spores packed within or surrounding the hair shaft (see Fig. 8.19e in Chapter 8). Hyphae are only rarely seen on the hair.

Scabies Preparation

A skin scraping showing a mite, its egg, or feces is necessary to diagnose infestation with *Acarus scabiei* because many other skin rashes resemble scabies clinically (Fig. 1.7). The most important factor for obtaining a successful scraping is the choice of site. Burrows and papules, which are most likely to harbor the mite, are commonly located on the wrists, fingers, and elbows. In infants, primary lesions may also be found on the trunk, palms, and soles. A fresh burrow can be identified as a 5–10 mm elongated papule, with a vesicle or pustule at one end. A small, dark spot resembling a fleck of pepper may be seen in the vesicle. This spot is the mite, and it can be lifted out of its burrow with a needle or the point of a scalpel. Usually, it is best to hold the skin taut between the thumb and index finger while vigorously scraping the burrow. Although this may induce a small amount of bleeding, if performed with multiple, short, rapid strokes, it is usually painless. A drop of mineral oil should be applied to the skin before scraping to ensure adherence of the scrapings to the blade. The scrapings are then placed on the slide, another drop of mineral oil is added, and a cover slip is applied. Gentle pressure with a pencil eraser may be used to flatten thick specimens.

Mites are eight-legged arachnids easily identified with the scanning power of the microscope. Care must be taken to focus through thick areas of skin scrapings so as not to miss any camouflaged mites. The presence of eggs (smooth ovals, approximately one half the size of an adult mite) or feces (brown pellets, often seen in clusters) are also diagnostic. If eggs or feces




Fig. 1.4 Primary skin lesions. Macule: a small (usually 1 cm), flat lesion showing an alteration in color or tone. Large macule is a patch. Papule: a small (1 cm), sharply circumscribed, elevated lesion. An elevated lesion over 1 cm is referred to as a plaque. Nodule: a soft or solid mass in the dermis or subcutaneous fat. Tumor: a large nodule, localized and palpable, of varied size and consistency. Vesicle: a blister containing transparent fluid. Bulla: a large blister. Wheal: an evanescent, edematous, circumscribed, elevated lesion that appears and disappears quickly. (Adapted from CIBA.)

are found first, perusal of the entire slide usually reveals the adult mite.

The dermatoscope can also be used to visualize the female mite whose mouth parts appear as an elongated triangle-shaped spot referred to as a delta sign.




Fig. 1.5 Secondary skin lesions. Scale: dry and/or greasy fragments of adherent epidermis. Pustule: a sharply circumscribed lesion containing free pus. Crust: a dry mass of exudate from erosions or ruptured vesicles/pustules, consisting of serum, dried blood, scales, and pus. Erosion: well-defined partial-thickness loss of epidermis. Ulcer: a clearly defined, full-thickness loss of epidermis that may extend into the subcutis. Scar: a permanent skin change resulting from new formation of connective tissue after destruction of the epidermis and cutis. When the loss of dermis and/or fat is prominent, the lesion may be atrophic. Fibrosis may result in firm, thickened papules or plaques. Excoriation: any scratch mark on the surface of the skin. Fissure: any linear crack in the skin, usually accompanied by inflammation and pain. (Adapted from CIBA.)

Lice Preparation

Lice are six-legged insects visible to the unaided eye that are commonly found on the scalp (Fig. 1.8), eyelashes, and pubic areas. Pubic lice are short and broad, with claws spaced far apart for grasping the sparse hairs on the trunk, pubic area, and




Fig. 1.6 Potassium hydroxide (KOH) preparation. **(a)** Small scales are scraped from the edge of the lesion onto a microscopic slide. **(b)** The scales are crushed to form a thin layer of cells in order to visualize the fungus easily. **(c)** In this positive KOH preparation of skin scrapings, fungal hyphae are seen as long septate, branching rods at the margins and center of the scales. **(d)** Pseudohyphae and spores typical of tinea versicolor give the appearance of spaghetti and meatballs.




Fig. 1.7 **(a)** Microscopic appearance of the adult scabies mite. Note the small oval egg within the body. **(b)** Scraping from an adolescent with crusted scabies shows two mites and multiple fecal pellets.

eyelashes, whereas scalp lice are long and thin, with claws closer together to grasp the denser hairs found on the head. The lice are best identified close to the skin, where their eggs are more numerous and more obvious. Diagnosis can be made by identifying the louse, or by plucking hairs and confirming the presence of its eggs or “nits” by microscopic examination.

A dermatoscope or magnifying glass can also be used for confirmation of lice infestation.

Tzanck Smear

The Tzanck smear is an important diagnostic tool in the evaluation of blistering diseases. It is most commonly used to




Fig. 1.8 Microscopic appearance of lice. **(a)** The crab louse has a short, broad body, with claws spaced far apart. **(b)** The head louse has a long, thin body, with claws closer together. **(c)** A hatched nit is tightly cemented to the hair shaft.

distinguish viral diseases, such as herpes simplex, varicella, and herpes zoster, from nonviral disorders (Fig. 1.9). It is important to note that Tzanck smears from vesicles of vaccinia and smallpox do not demonstrate multinucleated giant cells. The smear is obtained by removing the “roof” of the blister with a curved scalpel blade or scissors, and scraping the base to obtain the moist, cloudy debris. The material is then spread onto a glass slide, air dried, and stained with Giemsa or Wright stain. The diagnostic finding of viral blisters is the multinucleated giant cell. The giant cell is a syncytium of epidermal cells,




Fig. 1.9 Tzanck Smear. Note the multinucleated giant cells characteristic of viral infection with herpes simplex and varicella/zoster.

with multiple overlapping nuclei; it is much larger than other inflammatory cells. A giant cell may be mistaken for multiple epidermal cells piled on top of each other. If a microscope and stain are available, the Tzanck smear can be used for rapid confirmation of the clinical suspicion of infection, while more sensitive studies like polymerase chain reaction are pending.

Wood Light

Wood light is an ultraviolet source that emits at a wavelength of 365 nm. Formerly, its most common use was in screening patients with alopecia for tinea capitis, as the most common causative organisms, *Microsporum audouinii* and other *Microsporum* species, were easily identified by blue-green fluorescence under Wood light. However, today in North America, *Trichophyton* species are the most common fungi associated with tinea capitis, but it does not fluoresce. In the United States, fewer than 10% of cases are caused by *M. canis* and other *Microsporum* species. In Europe, Africa, and Asia, organisms that cause ectothrix scalp infection and which fluoresce include *M. ferrugineum*, *M. audouinii*, and *M. canis*.

Wood light is still of value in diagnosing a number of other diseases. Erythrasma is a superficial bacterial infection of moist skin in the groin, axilla, and toe webs. It appears as a brown or red flat plaque and is caused by a *Corynebacterium* that excretes a pigment that contains a porphyrin. This pigment fluoresces coral red or pink under Wood light. Tinea versicolor, a superficial fungal infection with hypopigmented macules and plaques on the trunk, also fluoresces under Wood light with a green-yellow color. *Pseudomonas* infection of the toe web space and colonization of the skin in burn patients will fluoresce yellow-green. Patients with porphyria cutanea tarda excrete uroporphyrins in their urine, and examination of a urine specimen will show an orange-yellow fluorescence. Adequate blood levels of tetracycline produce yellow fluorescence in the opening of hair follicles, while lack of fluorescence indicates poor intestinal absorption or poor patient compliance. Positive fluorescence in the skin is diagnostically useful, but many of the pigments that fluoresce are water soluble and readily removed by swimming or bathing.

Wood light also emits purple light in the visible spectrum. This wavelength can be used to accentuate subtle changes in pigmentation. The purple light is absorbed by melanin in the epidermis, and variably reflected by patches of hypopigmentation and depigmentation. It can be helpful to distinguish increased pigmentation in the epidermis that will be enhanced by the purple light from increased pigmentation in the dermis that does not enhance. Purple light may be particularly useful in evaluating light-pigmented individuals with vitiligo or ash leaf macules (congenital hypopigmented macules).

DERMATOLOGIC THERAPEUTICS

General Principles

Single component generic preparations are often effective and inexpensive. Fixed multiple component preparations are occasionally useful and may increase adherence to the treatment regimen, but the practitioner must be aware of all the constituent agents and the increased risk of adverse drug reactions. Specially formulated medications are often prohibitively expensive and seldom indicated in general practice. Fortunately, there are now some special formulating pharmacies that follow Food and Drug Administration guidelines, prepare safe and cost effective products, and obtain insurance preauthorization.

The practitioner must calculate the quantity of medication required for the patient to comply with instructions. In a child, 15–30 g of an ointment is needed to cover the entire skin surface once. This quantity will vary with the vehicle used and the experience of the individual applying the preparation.

Topical Vehicles

Two variables are particularly important in the selection of effective topical therapy: the active medication and the vehicle. No matter how effective the active medication, adherence to the recommended regimen will require that the clinician consider the selection of a vehicle carefully. In infants and young children, ointments tend to be better tolerated than other vehicles, while in older children, adolescents, and young adults, more elegant vehicles (e.g. creams, foams, sprays, solutions) that are free of lingering odors, color residues, or tackiness, will encourage adherence.

Ointments

In general, ointments are occlusive and allow for high transcutaneous penetration of the active drug. Ointments are stable for long periods and require few preservatives and bacteriostatic additives. As a consequence, they are least likely to cause contact allergy or irritation. These vehicles are well tolerated when the skin is cracked or fissured, particularly in young children with chronic skin disease (e.g. atopic dermatitis, psoriasis). Unfortunately, ointments tend to be messy and may stain clothing, so they are not often welcome by older children and adolescents.

Open Wet Dressings

Open wet dressings, using tap water or normal saline, provide symptomatic relief by cooling and drying acute inflammatory

lesions. They cleanse the skin by loosening exudates and crusts that can be painlessly removed before the dressing dries. Various astringents and antiseptics, such as vinegar or 5% aluminum acetate solution (e.g. Burrow solution), may be added to compression solutions in a 1:20–40 dilution. Bleach baths and chlorine swimming pools can also be used as gentle antiseptics and anti-inflammatory agents in patients prone to chronic or recurrent infection such as atopic dermatitis and epidermolysis bullosa. The concentration of chlorine in bleach baths (1/4 cup of household bleach or 59 mL to a 40-gallon bathtub to give a concentration of sodium hypochlorite of 6–8.25%) is designed to reproduce the concentration of chlorine recommended for supervised swimming pools.

Powders and Lotions

Powders promote drying and are especially useful in the intertriginous areas. Lotions are powders suspended in water (e.g. calamine lotion). When these preparations dry, they cool the skin and provide a uniform covering of the suspended agent. The clinician should warn patients and parents against the use of combination products that might result in irritation or percutaneous absorption of the active ingredients (e.g. calamine and diphenhydramine).

Gels

Gels are aqueous preparations that liquefy on contact with the skin and leave a uniform film on drying. Gels are well tolerated in hair-bearing areas. Water-based gels are best tolerated by children, while alcohol-containing products are more likely to cause burning or irritation. Gels are well tolerated in hair-bearing areas.

Aerosols

Aerosols and sprays act in a manner similar to lotions and gels. Active ingredients are incorporated into an aqueous phase. A convenient delivery system usually allows for easy dispersion over the skin surface. Aerosols are also particularly useful on the scalp.

Creams

Traditional creams are suspensions of oil in water. As the proportion of oil increases, the preparation approaches the consistency of an ointment, which is the most lubricating vehicle. Creams are water washable and hygroscopic. They may be drying and occasionally sensitizing.

Pastes

Pastes, which are mixtures of powder in ointment, are messy and may be difficult to remove from the skin. They are used to protect areas prone to irritation, such as the diaper area. Pastes can be removed with mineral oil.

Foams

Foams represent a novel vehicle, which enhances percutaneous absorption of medication in a cosmetically acceptable elegant

preparation. Foams remain stable until applied to the skin, where warming from natural body heat results in volatilization of inert contents with deposition of the active medication on the skin surface. Because foams contain minimal solid ingredients, there is little residue, making them particularly attractive vehicles for products designed for the scalp and intertriginous areas. A number of topical steroid foams have been approved for the treatment of atopic dermatitis and psoriasis, while other agents have been approved for seborrheic dermatitis and ichthyosis.

Shampoos and Washes

Short contact therapy with medicated shampoos and washes may also enhance adherence, particularly in adolescents with busy schedules and little time for topical therapy. These formulations contain insoluble particulate drugs such as benzoyl peroxide, salicylic acid, corticosteroids, and antifungal agents, some of which remain after showering or washing. Shampoos

and washes may also be particularly useful when longer periods of contact are likely to result in burning or irritation.

Topical Corticosteroids

Topical steroids are available in every type of vehicle. A good approach is to become familiar with one or two products in each of the potency ranges (Table 1.4). A check of local pharmacies is useful in determining the availability and cost of medications.

Most childhood skin eruptions requiring topical steroids can be readily managed with twice-daily applications of low- or medium-potency preparations. Moreover, a number of studies have shown that twice-daily applications of mid-potency agents can be applied to most areas of the skin in children for long periods of time safely. With few exceptions, low-potency medications should be used on the face and intertriginous areas because more potent preparations may produce atrophy, telangiectasias, and hypopigmentation. Regardless of

TABLE 1.4 Topical Corticosteroids

	Generic Name	Trade Name
Super high-potency topical steroids—class 1	Betamethasone dipropionate augmented 0.05%	Diprolene ointment 0.05%
	Clobetasol propionate 0.05%	Clobex lotion, spray, shampoo 0.05%
		Olux E foam 0.05%, Olux foam 0.05%
		Temovate cream, ointment, solution 0.05%
	Fluocinonide 0.1%	Vanos cream 0.01%
High-potency topical steroids—class 2	Halobetasol propionate 0.05%	Ultravate cream, ointment 0.05%
	Amcinonide 0.1%	
	Desoximetasone 0.25%, 0.05%	Topicort cream, ointment 0.25%, gel 0.05%
	Diflorasone diacetate 0.05%	ApexiCon E cream 0.05%
		Maxiflor ointment 0.05%
	Halcinonide 0.1%	Halog, Halog E ointment, cream 0.1%
Topical steroids—class 3	Fluocinonide 0.05%	Lidex cream, gel, ointment 0.05%
	Mometasone furoate 0.1%	Elocon ointment 0.1%
	Amcinonide 0.1%	Cyclocort cream, lotion 0.01%
	Betamethasone dipropionate 0.05%	Diprosone cream 0.05%
	Betamethasone valerate 0.1%	Valisone ointment 0.1%
		Betacap 0.1% (UK)
	Clobetasone butyrate 0.05%	Eumovate ointment, cream 0.05% (UK)
Topical steroids—class 4	Fluocinonide 0.05%	Lidex ointment, cream, gel 0.05%
	Fluticasone propionate 0.05%	Cutivate 0.05%, cream, lotion
	Betamethasone valerate 0.1%	Luxiq foam 0.1% ointment, cream, lotion
	Clocortolone pivalate 0.1%	Cloderm cream 0.1%
	Desoximetasone 0.05%	Topicort LP cream 0.05%
	Fluocinolone acetonide 0.025	Synalar ointment 0.025%
	Flurandrenolide 0.05%	Cordran ointment, lotion 0.05%
	Hydrocortisone probutate 0.1%	Pandel cream 0.1%
	Hydrocortisone valerate 0.2%	Westcort ointment 0.2%
	Prednicarbate 0.1%	Dermatop ointment 0.1%
	Triamcinolone acetonide 0.1%, 0.025%	Kenalog ointment 0.1%, 0.025%

TABLE 1.4 Topical Corticosteroids—cont'd

	Generic Name	Trade Name
Topical steroids—class 5	Betamethasone dipropionate 0.05%	Diprosone lotion 0.05%
	Betamethasone valerate 0.1%	Valisone cream, lotion 0.1%
	Fluocinolone acetonide 0.025%	Synalar 0.025%, cream 0.01%
	Fluticasone propionate 0.05%	Cutivate cream, lotion 0.05%
	Hydrocortisone butyrate 0.1%	Locoid Lipocream, ointment, lotion, solution 0.1%
	Hydrocortisone valerate 0.2%	Westcort cream 0.2%
	Prednicarbate 0.1%	Dermatop ointment, cream 0.1%
	Triamcinolone acetonide 0.1%	Kenalog cream, lotion 0.1%
Topical steroids—class 6	Alclometasone dipropionate 0.05%	Aclovate ointment, cream 0.05%
		Modrasone ointment, cream 0.05%
	Desonide 0.05%	DesOwen ointment, cream, lotion 0.05%
		Desonate Gel 0.05%
		Tridesilon cream 0.05%
	Fluocinolone acetonide 0.01%	Synalar cream, solution 0.1%
Topical steroids—class 7		Derma-Smoothe/FS oil
	Hydrocortisone 2.5%	Hytone cream, lotion 2.5%
		Cobadex 1%
		Dioderm 0.1%
		Mildison 1%
		Hydrocortisyl 1%
		Hytone ointment 1%
	Dexamethasone 0.04%	Hexadrol cream 0.04%
	Methylprednisolone acetate 0.25%	Medrol ointment 0.25%
Topical steroids—class 8	Prednisolone 0.5%	Meti-derm cream 0.5%
	Hydrocortisone 0.5%, 1%	Hytone, Cortaid, Synacort, Nutracort 0.5% ointment, cream, lotion

the potency of a medication, patients should be followed carefully for steroid-induced changes, even though they are only rarely produced by therapy restricted to 2–4 weeks. Patients receiving chronic therapy to sensitive areas should take frequent “time-outs” from their topical steroids (e.g. 1 week per month) and should taper them when possible. Tapering may be achieved by decreasing the frequency of application, as well as by mixing the active preparation with a bland emollient such as petrolatum.

Steroids may mask infections and suppress local and systemic immune responses. Consequently, they are contraindicated in most patients with viral, fungal, bacterial, or mycobacterial infections.

Topical Calcineurin Inhibitors

The topical nonsteroidal anti-inflammatory agents, pimecrolimus (Elidel) and tacrolimus (Protopic), provide an alternative for the treatment of atopic dermatitis. These calcineurin inhibitors selectively suppress the release of inflammatory mediators from lymphocytes without compromising the function of melanocytes, fibroblasts, or endothelial cells. As a consequence, they are not associated with the development of pigment alteration, atrophy, or telangiectasias. They can be applied at any site including the genital skin, breasts, and face. However,

they are contraindicated in erythrodermic conditions, where significant percutaneous absorption may occur.

The nonsteroidal agents are approved for use in children over 2 years of age, but recent studies on a large number of patients from 3 months to 2 years old demonstrate safety and efficacy similar to older children.

A black box warning cautions prescribers and patients against using these agents in children under 2 years of age and long term in any patient. However, when used judiciously, they offer a safe alternative to topical steroids particularly in sensitive areas of the skin. Before using these agents, it is imperative that the practitioner discuss the black box warning and the rationale for prescribing them.

Emollients (Lubricants)

Another topical non-steroidal agent crisaborole ointment 2%, which is anti-inflammatory phosphodiesterase-4 inhibitor, is safe and approved for children with eczema down to 3 months of age. Any preparation that reduces friction and leaves a smooth, occlusive film that prevents drying is classified as a lubricant (Table 1.5). In patients with chronic dermatitis, ointments (or water-in-oil-based products) provide the best lubrication, especially during the dry winter months. Less oily preparations (oil-in-water creams, lotions, foams, aerosols) are

TABLE 1.5 Emollients

Types of Skin	Moisturizing Base Type	Product Name
1. Extremely dry skin	Ointment or oil-based	Bag Balm
		Blue Star ointment
		Elta Swiss skin cream
		Johnson's Baby Oil
		Palmer's Cocoa Butter
		Theraplex Emollient
		Vaseline Petroleum Jelly
		Mineral Oil
		A+D ointment
		Alpha Keri Moisture Rich Baby Oil
		Aquaphor Healing ointment
2. Dry	Water-in-oil emulsion	A+D ointment with zinc oxide
		Acid Mantle cream
		Elta Light moisturizing cream
		Eucerin Original moisturizing cream/lotion
		Jergens All-Purpose Face Cream
		Olay Body lotion
		Restoraderm lotion
		St. Ives Swiss Formula products
		Sween Cream
		Theraplex Clear lotion
		Vanicream
		Vaseline Intensive Care lotion
3. Normal to dry	Oil-in-water	Alpha hydroxy cream/lotion
		Aqua Care cream
		Biore Balancing Moisturizer Normal to Dry
		Caress Body Silkening lotion
		Carmol 40 cream
		Complex 15 lotion
		Curél Moisturizing lotion
		Cutemol cream
		Gold Bond Moisturizing Body Lotion
		Jergens Original Scent lotion
		Keri lotion
		LactiCare lotion
		Lubriderm Skin Therapy
		Moisturel cream/lotion
		Neutrogena lotion
		Nivea Body Creamy Conditioning Oil
		Nutraderm lotion
		Olay Active Hydrating Original Cream
		Pacquin Plus skin cream
		Pond's Age Defying lotion/cream
		Purpose Alpha Hydroxy Moisture cream/lotion
		Sarna lotion
4. Normal to oily	Oil-free	Carmol 10 Deep Moisturizing lotion
		CeraVe lotion or cream
		Cetaphil lotion or cream
		Corn Huskers lotion
		Epilyt lotion
		Gerber Baby Lotion
		Johnson's Baby Lotion
		Lubriderm Skin Therapy
		Neutrogena Combination Skin Moisture
		Olay Regenerist Facial Moisturizer
		Walgreens Glycerin and Rosewater

often preferred by patients during the spring and summer. More elegant products should be considered in older children and young adults, especially for use on the scalp or intertriginous areas. Cultural preferences should also be taken into account when selecting a lubricant. Preparations containing topical sensitizers such as fragrance, neomycin, and benzocaine should be avoided, particularly in patients with inflamed skin.

Sun Protective Agents

These agents (Table 1.6) include sunscreens (light-absorbing compounds) and sunblocks (inert compounds that reflect light). Although the long-awaited guidelines from the Food and Drug Administration have not yet been finalized, in 2019 the Food and Drug Administration recently described two sun-blocking ingredients, titanium dioxide and zinc oxide, as generally recognized as safe and effective (GRASE). The Food and Drug Administration also listed two other ingredients para-aminobenzoic acid (PABA) and trolamine salicylate as not GRASE and 12 other ingredients as potentially safe ingredients pending the presentation of more data on safety and efficacy (see Table 1.6 for listing of GRASE and potentially GRASE ingredients). Most experts recommend the use of broad-spectrum sun protective agents (protective against both ultraviolet A and B light) with an SPF of at least 30. Parents should also be counseled to purchase products that are water resistant. See Chapter 7 Photodermatoses for further discussion of these agents.

TABLE 1.6 Sun Protection

Sunlight Type	Ultraviolet Spectrum (nm)	Sun Protection Agent
Visible light	>400	Titanium dioxide
		Zinc oxide
UVA	320–400	Dioxybenzone
		Titanium dioxide
		Zinc oxide
		Avobenzone
		Octocrylene
		Avobenzone
		Meradimate
		Cinoxate
		Titanium dioxide
		Zinc oxide
		Oxybenzone
		Sulisobenzonate
		Octocrylene
UVB	290–320	Cinoxate
		Padimate
		Homosalate
		Octisalate
		Ensulizole
		Filtered out by ozone, does not reach the Earth's surface
UVC	<290	Filtered out by ozone, does not reach the Earth's surface

UVA, Ultraviolet light A; UVB, ultraviolet light B; UVC, ultraviolet light C.

FURTHER READING

- Bologna JL, Schaeffer JV, Cerroni L. Dermatology, 4th edn. Elsevier, China, 2018.
- Baran R, de Berker D, Holzberg M, et al. Diseases of the nails and their management, 5th edn. Wiley & Sons, Ltd, India, 2019.
- Cragi FE, Smith E, Williams H. Bleach baths to reduce severity of atopic dermatitis colonized by staphylococcus. Arch Dermatol 2010;146:541-543.
- Eichenfield LF, Frieden IJ. Neonatal and infant dermatology, 3rd edn. Elsevier, Saunders, China, 2015.
- Elder DE, Elenitsas R, Rosenbach M, et al. Lever's Histopathology of the skin, 11th edn. Lippincott, Wolters Kluwer, China, 2015.
- Habif TP. Clinical dermatology: a color guide to diagnosis and therapy, 6th edn. Saunders, St Louis, 2016.
- Harper JH, Oranje A, Prose N. Harper's textbook of pediatric dermatology, 3rd edn. Wiley-Blackwell Science, London, 2011.
- Hordinsky MK, Sawaya ME, Scher RK. Atlas of hair and nails. Churchill Livingstone, Philadelphia, 2000.
- Kang S, Amagai M, Bruckner AL, et al. Fitzpatrick's dermatology, 9th edn. McGraw Hill, New York, 2019.
- Kristal L, Prose N. Color atlas of pediatric dermatology, 5th edn. McGraw Hill, China, 2017.
- Neville BW, Damm DD, Allen CM, Chi AC. Oral and maxillofacial pathology, 4th edn. Elsevier, Canada, 2016.

- Paller AS, Mancini AJ. Hurwitz clinical pediatric dermatology, 5th edn. Elsevier, Canada, 2016.
- Patterson JW. Weedon's skin pathology, 4th edn. Churchill Livingstone, Elsevier, China, 2016.
- Schachner LA, Hansen RC. Pediatric dermatology, 4th edn. CV Mosby, New York, 2011.

USEFUL WEBSITES

- A good resource for skin diseases, conditions, and treatments: <https://www.dermnetnz.org>
- General medical reference with data from a number of medical texts: <https://www.emedicine.com>
- National Library of Medicine: <https://www.nlm.nih.gov>
- PubMed: National Library of Medicine reference journal database: <https://www.ncbi.nlm.nih.gov/pubmed>
- Online Mendelian Inheritance in Man: <https://www.ncbi.nlm.nih.gov/omim>
- A good resource for pediatricians and families: <https://www.aad.org/public/kids>
- See section for patients and families: <https://pedsderm.net>